



Management of Spinal Lymphomas: Spinal Instability Assessment Using the Spinal Instability Neoplastic Score and a Proposed Treatment Algorithm

Ali Fatih RAMAZANOGLU¹, Mustafa Umut ETLI¹, Caner SARIKAYA¹, Mehmet Resid ONEN², Ali Hakan KAYA³, Sait NADERI¹

¹Health Sciences University, Umraniye Training and Research Hospital, Department of Neurosurgery, Istanbul, Turkey

²VM Medical Park Maltepe Hospital, Department of Neurosurgery, Istanbul, Turkey

³Health Sciences University, Umraniye Training and Research Hospital, Department of Hematology, Istanbul, Turkey

Corresponding author: Mustafa Umut ETLI ✉ umutetli@gmail.com

ABSTRACT

AIM: To propose a treatment algorithm, and to assess spinal instability in patients diagnosed with spinal lymphoma.

MATERIAL and METHODS: Demographics, symptoms, tumor level and location, and presence of spinal instability were reviewed in 22 patients with spinal lymphomas. Each patient's neurological state was reviewed using the American Spinal Injury Association and modified McCormick scale scores, and spinal instability was assessed using the Spinal Instability Neoplastic Score (SINS).

RESULTS: Initially, percutaneous biopsy was performed in 16 patients, and open biopsy was performed in 6 patients. Eight of the patients who underwent percutaneous biopsy were followed up with hematological examination alone, as they had no additional complaints. The SINS was used to evaluate the presence of spinal instability, and the type of surgery to be performed was decided accordingly. In the second surgery, decompression and stabilization were performed in 5 of the remaining 8 patients, and only decompression was performed in 3 of them. Neurological improvement was observed in 6 of 7 patients with acute neurological deficit.

CONCLUSION: Percutaneous biopsy for tissue diagnosis is the first step in the management of spinal lymphomas. Patients without neurological deficit should be referred for hematological examination. Those with acute neurological deficit require emergency surgery, and those with chronic symptoms must undergo operation for decompression and/or stabilization. This study confirmed the safety of the SINS in the evaluation of spinal instability in spinal lymphoma cases.

KEYWORDS: Management, SINS, Spinal instability, Spinal lymphoma

ABBREVIATIONS: **ASIA:** The American Spinal Injury Association; **CT:** Computed tomography, **MR:** Magnetic resonance, **NOMS:** Neurologic, oncologic, mechanical and systemic decision framework, **OB:** Open biopsy, **PB:** Percutaneous biopsy, **SINS:** The Spinal Instability Neoplastic Score, **SL:** Spinal lymphoma

INTRODUCTION

Spinal lymphomas (SLs) constitute 9%–10% of all epidural malignancies (2,4,17). They are localized in the paraspinal, vertebral, and epidural areas (13,15,19,25).

Spinal cord compression due to SL is extremely rare and is found in approximately 0.1%–3.3% of SL cases (15,26). Although medical treatment is effective for most cases of SL, some cases may require surgical intervention for pain relief,

Ali Fatih RAMAZANOGLU  : 0000-0001-6500-5070
Mustafa Umut ETLI  : 0000-0003-0756-4274
Caner SARIKAYA  : 0000-0002-2325-3903

Mehmet Resid ONEN  : 0000-0003-4353-9068
Ali Hakan KAYA  : 0000-0003-0496-0984
Sait NADERI  : 0000-0002-6784-4270

decompression, and/or spinal reconstruction (1,3,10,11,16–18,20,21,23,32,34).

On the other hand, assessment of spinal instability is still controversial in patients with SL (7,16,18,21,28,31). The available literature indicates that the Spinal Instability Neoplastic Score (SINS) has the potential to be a valuable guide for the management of patients with spinal metastases (9,24). The purpose of this study was to review the treatment results of SL cases, present our treatment algorithm, and report the reliability of the SINS for the assessment of SLs.

MATERIAL and METHODS

Twenty-two patients who underwent surgery for SL in our clinic between January, 2011 and December, 2019 were retrospectively evaluated. The patient population consisted of 14 men and 8 women, with a mean age of 34.2 years (range: 15–81 years).

In this study, demographic characteristics, symptomatology (pain and neurological findings), level with respect to the spine segments, location with respect to the dura, biopsy results, and surgical procedures were reviewed. Patients who underwent an operation for SL in our clinic between January, 2011 and December, 2019 were included in the study. Radiological diagnoses were made using magnetic resonance (MR) imaging, computed tomography (CT), and positron emission tomography-CT. Furthermore, CBC, peripheral blood smear or bone marrow aspiration were performed in elective cases for differential diagnosis. The American Spinal Injury Association (ASIA) and modified McCormick scales were used to evaluate preoperative and postoperative neurological function (Table I), and the SINS score was used to evaluate the presence of spinal instability (5,8). Except for the patients who presented with neurological deficit, the treatments to be applied in the follow-up were decided together with the oncology council. At the decision of the oncology team, biopsy was performed both for staging and excluding a possible second metastatic tumor. Percutaneous biopsy specimens were obtained by Jamshidi needle and biopsy punch through the pedicle into the site of the disease using C-arm fluoroscopy.

RESULTS

The tumors were located in the lumbar spine in 15 patients, in the thoracic spine in 5 patients, and in the sacral spine in

2 patients. The vertebral body was involved in all the cases. Whereas the tumors were purely vertebral in 2 cases, extension was observed to the paraspinal area in 3 cases, to the epidural area in 2 cases, to the paraspinal and epidural areas in 6 cases, to the prevertebral and epidural areas in 1 case, and to the prevertebral, vertebral, epidural, and paraspinal areas in 8 cases. Only 2 patients had a hematologically confirmed SL before diagnosis, and 20 patients were admitted with an undiagnosed spinal tumor. Seventeen patients were admitted with spine-related pain, and 7 patients had neurological complaints, such as neuropathic pain, numbness, and neurological deficit.

In the preoperative neurological evaluation, the neurological function was ASIA grade B in 2 patients, ASIA grade C in 3 patients, ASIA grade D in 4 patients, and ASIA grade E in 13 patients. In the postoperative neurological evaluation, 1 patient had ASIA grade C, 5 patients had ASIA grade D, and 16 patients had ASIA grade E neurological function.

The preoperative modified McCormick scale score was grade 1 in 15 patients, grade 2 in 2 patients, grade 3 in 3 patients, and grade 4 in 2 patients. Grade 1 became grade 3 in 1 patient soon after percutaneous biopsy (PB), so decompression surgery was performed (Patient no. 18). The postoperative modified McCormick evaluation results were as follows: grade 1 in 17 patients, grade 2 in 3 patients, and grade 3 in 2 patients. The preoperative SINS score was 6 points in 1 patient, 7–12 points (potentially unstable) in 14 patients, and >12 points (unstable) in 7 patients.

A PB was performed in 16 patients, and open biopsy (OB) was performed in 6 patients. Of the 16 patients who underwent PB, 8 who had neither severe pain nor neurological deficit were referred to the hematology department for further treatment of lymphoma (Figure 1). 13 patients with ASIA E were diagnosed by PB to confirm the pathological diagnosis. Of these 13 patients' 6 were operated because of potential instability (high SINS score), and two sacral lesions for relieving urinary retention of acute onset (patient no. 12 and 22).

A decompression procedure was performed in 3 of the 16 patients who underwent PB (Figure 2), and stabilization and decompression were performed in the remaining 5 cases (Figure 3). OB of 6 patients was performed during decompression procedure. 4 of these cases underwent only decompression, and 2 underwent stabilization and decompression. In total,

Table I: Preoperative and Postoperative Modified McCormick Scale Scores

Total no of patients = 22	Postoperative Modified McCormick Scale				
	I	II	III	IV	V
I	14				
II	2				
III		2	2		
IV	1	1			
V					



Figure 1: Sagittal T2 weighted (A) and post-contrast T1 weighted MR images (B) of a case with low back pain showed presence of tumor at the level of L1-2. PB confirmed presence of lymphoma and patient referred to the hematology. Five years later MR images showed no tumor at sagittal T2 weighted (C) and post-contrast T1 weighted MR images (D).



Figure 2: Sagittal T2 weighted (A), sagittal postcontrast T1 weighted (B) and axial postcontrast T1 weighted (C) MR images of a case with low back pain and presence of tumor compressing the spinal cord at the level of sacrum. Decompression surgery performed after PB. One year later MR images showed decrease in compression at sagittal T2 weighted (D), sagittal post-contrast T1 weighted (E) and axial post contrast T1 weighted (F) MR images.




Figure 3: Sagittal T2 weighted (A), sagittal postcontrast T1 weighted (B) and axial postcontrast T1 weighted (C) MR images of a case with low back pain and progressive paraparesis showed presence of tumor compressing the spinal cord circumferentially at level of T11-12. Open biopsy, decompression and stabilization performed to the patient. The patient improved neurologically and referred to the hematology. One year later, sagittal T2 weighted (D), sagittal post-contrast T1 weighted (E) and axial post contrast T1 weighted (F) MR images showed no tumor.

7 patients underwent decompression (3 after PB and 4 after OB), and 7 patients underwent stabilization and decompression (5 after PB and 2 after OP).

Neurological improvement was observed in 6 of the 7 patients with acute neurological deficit.

DISCUSSION

This study demonstrated that PB for tissue diagnosis should be the first step in the management of SL with or without minor symptoms. Patients with major symptoms, such as severe pain and neurological deficit, may require OB with decompression and/or stabilization. Although there is a known pathological diagnosis in patients with spinal metastases, the possibility of a second pathological origin should be considered for synchronous or metachronous tumor (27).

Evaluation of spinal instability for non-spine surgeon clinicians is not easy. This study demonstrates the reliability of the SINS in evaluating spinal instability in SLs. Neurologic, oncologic, mechanical, and systemic (NOMS) decision framework is a classification that has been used for more than two decades and examines spinal metastases in detail (14). The SINS has been used for evaluation of M section (mechanical) of NOMS.

According to the World Health Organization, lymphomas can be basically classified as Hodgkin and non-Hodgkin lymphomas (29). SLs, with or without extravertebral extension,

should be classified as bone lymphomas. SLs are commonly non-Hodgkin lymphomas, with a predominance of diffuse large B-cell lymphomas (50%-80%) (33). The cases presented here are non-Hodgkin lymphomas.

Most (90%) spinal tumor cases treated surgically in our clinic are metastatic cancers, followed by hematological malignancies such as multiple myeloma, plasmacytoma, and SL. SLs account for 40% (22 cases) of the 55 hematological malignancies involving the spine in our spine tumor series, and epidural extension was detected in 68% of the 22 cases.

The management of SL is still controversial (Table II). Whereas some authors recommend medical treatment, others recommend surgery for decompression and stabilization. Another controversy concerns the assessment of spinal instability, which is relatively difficult for non-spine surgeons.

The effectiveness of chemotherapy and radiotherapy has been reported by many authors (1,23,32,34). As in other neoplastic spinal cord compressions, the presence of progressive acute neurologic deficit and/or spinal instability indicates the need for acute surgery for biopsy, decompression, and stabilization. The importance of early surgery in neurologically involved SL cases was reported by Han et al. in 37 cases and by Chang et al. in 11 cases (3,11). Acute surgery was performed in 7 of the 22 cases in the present series, all except 1 of which improved significantly. Therefore, we agree with the aforementioned

Table II: Level, Localization, SINS, Type of Surgery, and Pathological Results of the Cases

Patient no	Tumor Level	Localization	SINS	1 st surgery	2 nd surgery	Pathological Diagnosis	Preoperative Neurological Examination
1	T10	PV+ V + E + PS	14	PB	D + I	Anaplastic large T cell lymphoma	McCormick 1; Unable to walk associated with dysesthesia.
2	L2-3	PV+ V + E	13	PB	D + I	High-grade B cell	McCormick 1
3	L5	V + PS	6	PB		High-grade B cell	McCormick 1
4	L5	PV+ V + E + PS	10	PB		B cell	McCormick 1
5	L1	PV+ V + E + PS	10	PB		T cell	McCormick 1
6	T 11-12	V + E + PS	13	PB	D + I	Non-Hodgkin	McCormick 3 (chronic)
7	L4	V + E + PS	14	D + I		Large B cell	McCormick 4
8	L4	V + PS	9	PB		Low-grade B cell	McCormick 1
9	T11-L3	V + PS	9	D		High-grade B cell	McCormick 3
10	L1-2	V + E + PS	10	PB		B cell	McCormick 1
11	C6-T1	V + E	10	D		B-cell prediagnosis	McCormick 4
12	Sacrum	PV+ V + E + PS	9	PB	D	B cell	McCormick 1
13	L5	V	10	PB		B cell	McCormick 1
14	L5	PV+ V + E + PS	15	PB	D + I	B cell	McCormick 1
15	L5	PV+ V + E + PS	14	PB	D + I	Non-Hodgkin	McCormick 1
16	L1	V	8	PB		B cell	McCormick 1
17	L1-4	PV+ V + E + PS	8	D		Diffuse large B cell	McCormick 2
18	T6	PV+ V + E + PS	10	PB	D	Diffuse large B cell	McCormick 1; 1 week after PB McCormick 3
19	L5	V + E + PS	10	PB		B-cell prediagnosis	McCormick 1
20	L5-Sacrum	V + E	8	D		Low-grade B cell	McCormick 2
21	L2-3-4	V + E + PS	13	D + I		Diffuse large B Cell	McCormick 3
22	Sacrum	V + E + PS	9	PB	D	Plasmablastic lymphoma	McCormick 1

E: Epidural, V: Vertebral, PV: Prevertebral, PS: Paraspinal, PB: Percutaneous biopsy, D: Decompression, I: Instrumentation.

authors on the importance and effectiveness of emergency surgery in SL cases with acute neurological deficit. Neurological deficit in such cases is not only due to epidural tumor compression. Spinal instability and kyphotic changes also cause neurological deficit, as well as pain. Therefore, clinicians should evaluate neoplastic spinal instability. SINS system is effective and provides easy assessment of neoplastic spinal instability. It was first introduced for the assessment of instability in spinal metastasis, particularly by non-surgeon clinicians (6). Its reliability and effectiveness in spinal metastasis have been reported by many authors (5,6,8,22). To the best of our knowledge, this is the first study to address the assessment of spinal instability in SL using SINS. Our study demonstrated that patients with scores > 12 have potential or overt neoplastic instability and are candidates for stabilizing surgery.

This study introduces the treatment algorithm we developed for patients with SL (Figure 4). On the basis of the results of 22 SL cases, we recommend PB for patients with or without minor symptoms. PB should be performed to confirm the diagnosis of SL and exclude other primary and metastatic tumors. These cases are referred to the hematology department for hematological treatment, including chemotherapy and radiotherapy. Although some reports on the use of vertebroplasty in the treatment of SLs, no vertebroplasty was performed in the present series (12,30).

The limitations of the study are that it is retrospective and the number of patients is small, however, larger case series are needed in future studies.

Patients with acute progressive deficit should undergo operation as soon as possible, as mentioned previously.




Figure 4: Proposed treatment algorithm for spinal lymphomas.

Patients with spinal instability and chronic deficit should undergo operation for elective decompression and stabilization after PB. This group requires a multidisciplinary approach for determining the appropriate timing of surgery.

CONCLUSION

We conclude that patients with SL who have minor neurological deficit require PB for tissue diagnosis. The presence of chronic neurological deficit and spinal instability may require surgery for elective decompression and stabilization. Patients with acute deficit due to tumor or instability-related compression require emergency decompression and stabilization. The use of SINS system facilitates the assessment of spinal instability by non-surgeon clinicians.

ACKNOWLEDGEMENT

Preparation for publication of this article is partly supported by Turkish Neurosurgical Society.

AUTHORSHIP CONTRIBUTION

Study conception and design: AFR, MUE, SN

Data collection: CS, MUE

Draft manuscript preparation: AFR, MUE

Critical revision of the article: SN

Other (study supervision, fundings, materials, etc.): AHK, MRO

All authors (AFR, MUE, CS, MRO, AHK, SN) reviewed the results and approved the final version of the manuscript.

REFERENCES

1. Aabo K, Walbon-Jørgensen S: Central nervous system complications by malignant lymphomas: Radiation schedule and treatment results. *Int J Radiat Oncol Biol Phys* 12:197-202, 1986
2. Boukobza M, Mazel C, Touboul E: Primary vertebral and spinal epidural non-Hodgkin's lymphoma with spinal cord compression. *Neuroradiology* 38:333-337, 1996
3. Chang CM, Chen HC, Yang Y, Wang RC, Hwang WL, Teng CLJ: Surgical decompression improves recovery from neurological deficit and may provide a survival benefit in patients with diffuse large B-cell lymphoma-associated spinal cord compression: A case-series study. *World J Surg Oncol* 11:90, 2013
4. Cugati G, Singh M, Pande A, Ramamurthi R, Balasubramanyam M, Sethi S, Singh A: Primary spinal epidural lymphomas. *J Craniovertebr Junction Spine* 2:3-11, 2011

5. Fisher CG, Dipaola CP, Ryken TC, Bilsky MH, Shaffrey CI, Berven SH, Harrop JS, Fehlings MG, Boriani S, Chou D, Schmidt MH, Polly DW, Biagini R, Burch S, Dekutoski MB, Ganju A, Gerszten PC, Gokaslan ZL, Groff MW, Liebsch NJ, Mendel E, Okuno SH, Patel S, Rhines LD, Rose PS, Sciubba DM, Sundaresan N, Tomita K, Varga PP, Vialle LR, Vrionis FD, Yamada Y, Fourny DR: A novel classification system for spinal instability in neoplastic disease: An evidence-based approach and expert consensus from the Spine Oncology Study Group. *Spine (Phila. Pa. 1976)* 35(22):E1221-9, 2010
6. Fisher CG, Schouten R, Versteeg AL, Boriani S, Varga PP, Rhines LD, Kawahara N, Fourny D, Weir L, Reynolds JJ, Sahgal A, Fehlings MG, Gokaslan ZL: Reliability of the Spinal Instability Neoplastic Score (SINS) among radiation oncologists: An assessment of instability secondary to spinal metastases. *Radiat Oncol* 9:69, 2014
7. Flouzat-Lachaniette CH, Allain J, Roudot-Thoraval F, Poinard A: Treatment of spinal epidural compression due to hematological malignancies: A single institution's retrospective experience. *Eur Spine J* 22:548-555, 2013
8. Fourny DR, Frangou EM, Ryken TC, DiPaola CP, Shaffrey CI, Berven SH, Bilsky MH, Harrop JS, Fehlings MG, Boriani S, Chou D, Schmidt MH, Polly DW, Biagini R, Burch S, Dekutoski MB, Ganju A, Gerszten PC, Gokaslan ZL, Groff MW, Liebsch NJ, Mendel E, Okuno SH, Patel S, Rhines LD, Rose PS, Sciubba DM, Sundaresan N, Tomita K, Varga PP, Vialle LR, Vrionis FD, Yamada Y, Fisher CG: Spinal instability neoplastic score: An analysis of reliability and validity from the spine Oncology Study Group. *J Clin Oncol* 29:3072-3077, 2011
9. Fox S, Spiess M, Hnenny L, Fourny DR: Spinal instability neoplastic score (SINS): Reliability among spine fellows and resident physicians in orthopedic surgery and neurosurgery. *Glob Spine J* 7:744-748, 2017
10. Haddad P, Thael JF, Kiely JM, Harrison EG, Miller RH: Lymphoma of the spinal extradural space. *Cancer* 38:1862-1866, 1976
11. Han S, Yang X, Jiang D, Zhou W, Liu T, Yan W, Wang T, Cheng L, Xiao J: Surgical outcomes and prognostic factors in patients with diffuse large b-cell lymphoma-associated metastatic spinal cord compression. *Spine (Phila. Pa. 1976)* 41:E943-E948, 2016
12. Jensen ME, Kallmes DF: Percutaneous vertebroplasty in the treatment of malignant spine disease. *Cancer J* 8:194-206, 2002
13. Kapoor R, Kumar V, Sharma SC: Primary extradural non-hodgkin's lymphoma. *JK Science* 8(1):January-March, 2006
14. Laufer I, Rubin DG, Lis E, Cox BW, Stubblefield MD, Yamada Y, Bilsky MH: The NOMS framework: Approach to the treatment of spinal metastatic tumors. *Oncologist* 18:744, 2013
15. Lim CC, Chong BK: Spinal epidural non-Hodgkin's lymphoma: Case reports of three patients presenting with spinal cord compression. *Singapore Med J* 37:497-500, 1996
16. Lyons MK, O'Neill BP, Kurtin PJ, Richard Marsh W: Diagnosis and management of primary spinal epidural non-Hodgkin's lymphoma. *Mayo Clin Proc* 71:453-457, 1996
17. Mally R, Sharma M, Khan S, Velho V: Primary lumbo-sacral spinal epidural non-Hodgkin's lymphoma: A case report and review of literature. *Asian Spine J* 5:192-195, 2011
18. McDonald AC, Nicoll JAR, Rampling RP: Non-Hodgkin's lymphoma presenting with spinal cord compression; a clinicopathological review of 25 cases. *Eur J Cancer* 36:207-213, 2000
19. McLain RF, Markman M, Bukowski RM, Macklis R, Benzel EC: *Cancer in the Spine: Comprehensive Care (Current Clinical Oncology)* 2006th ed. Humana, 2008:1-382
20. Monnard V, Sun A, Epelbaum R, Poortmans P, Miller RC, Verschueren T, Scandolaro L, Villa S, Majno SB, Ostermann S, Ozsahin M, Mirimanoff RO: Primary spinal epidural lymphoma: Patients' profile, outcome, and prognostic factors: A multicenter Rare Cancer Network study. *Int J Radiat Oncol Biol Phys* 65:817-823, 2006
21. Moussaly E, Nazha B, Zaarour M, Atallah JP: Primary non-hodgkin's lymphoma of the spine: A case report and literature review. *World J Oncol* 6:459-463, 2015
22. Murtaza H, Sullivan CW: Classifications in brief: The spinal instability neoplastic score. *Clin Orthop Relat Res* 477:2798-2803, 2019
23. Peng X, Wan Y, Chen Y, Chen L, He A, Liao W, Shen J, Fu Q, Han S, Li F, Zou X: Primary non-Hodgkin's lymphoma of the spine with neurologic compression treated by radiotherapy and chemotherapy alone or combined with surgical decompression. *Oncol Rep* 21:1269-1275, 2009
24. Pennington Z, Ahmed AK, Cottrill E, Westbroek EM, Goodwin ML, Sciubba DM: Intra- and interobserver reliability of the Spinal Instability Neoplastic Score system for instability in spine metastases: A systematic review and meta-analysis. *Ann Transl Med* 7:218-218, 2019
25. Rubinstein LJ: *Tumors of the Central Nervous System*. Washington DC, 1972. Available from: https://books.google.com/books?hl=tr&lr=&id=xKWxCqzfN5sC&oi=fnd&pg=PA1&d-q=Rubinstein+LJ.+Tumor+of+the+nervous+system&ots=i2E-0jyb1CU&sig=7ekwl_y77Ny6ZxwINFlxq2MWBqo
26. Salvati M, Cervoni L, Artico M, Raco A, Ciappetta P, Delfini R: Primary spinal epidural non-Hodgkin's lymphomas: A clinical study. *Surg Neurol* 46:339-343, 1996
27. Sohn S, Chung CK, Jung JH, Lee KC, Kim J, Chang UK, Sohn MJ, Kim SH: Nationwide comparative study of synchronous and metachronous spine metastasis in the adult Korean population. *J Clin Neurosci* 57:33-37, 2018
28. Spinazzé S, Caraceni A, Schrijvers D: Epidural spinal cord compression. *Crit Rev Oncol Hematol* 56:397-406, 2005
29. Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, Advani R, Ghielmini M, Salles GA, Zelenetz AD, Jaffe ES: The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood* 127(20):2375-2390, 2016
30. Tang H, Jia P, Li J, Chen H, Bao L, Feng F: Literature review percutaneous vertebroplasty for primary non-hodgkin's lymphoma of the thoracic spine: Case report and review of the literature case report. *Pain Physician* 20:727-735, 2017
31. Tang Y, Yang X, Xiao J, Liu K, Yan W, Song D, Qu J, Ma J, Wan W: Clinical outcomes of treatment for spinal cord compression due to primary non-Hodgkin lymphoma. *Spine J* 13:641-650, 2013

32. Wada N, Kohara M, Ikeda J, Hori Y, Fujita S, Okada M, Ogawa H, Sugiyama H, Fukuhara S, Kanamaru A, Hino M, Kanakura Y, Morii E, Aozasa K: Diffuse large B-cell lymphoma in the spinal epidural space: A study of the Osaka Lymphoma Study Group. *Pathol Res Pract* 206:439-444, 2010
33. Wang Y, Li J, Wei R, Liu C, Nataraj A, Yan J: Prognostic factors associated with bone lymphoma primarily presenting in the spine. *Spine (Phila. Pa. 1976)* 44:185-194, 2019
34. Xiong L, Liao LM, Ding JW, Zhang ZL, Liu AW, Huang L: Clinicopathologic characteristics and prognostic factors for primary spinal epidural lymphoma: Report on 36 Chinese patients and review of the literature. *BMC Cancer* 17:131, 2017